

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 January 2002 (17.01.2002)

PCT

(10) International Publication Number
WO 02/03915 A2

(51) International Patent Classification⁷: **A61K**

(21) International Application Number: PCT/US01/21509

(22) International Filing Date: 9 July 2001 (09.07.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/613,692 10 July 2000 (10.07.2000) US

(71) Applicant and

(72) Inventor: **KOZACHUK, Walter, E.** [US/US]; 11403-B
Cam Court, Kensington, MD 20895-1313 (US).

(74) Agent: **LINIAK, Thomas, P.**; Liniak, Berenato, Longacre & White, LLC, Suite 240, 6550 Rock Spring Drive, Bethesda, MD 20817 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHODS OF PRODUCING WEIGHT LOSS AND TREATMENT OF OBESITY

(57) Abstract: Methods are disclosed for the acute and chronic treatment of obesity using drugs whose mechanism includes the interaction and antagonism of the kainate/AMPA receptor. Methods are disclosed for employing the drug topiramate (topomax) as monotherapy or in combination therapy with lamotrigine, valproic acid, valproic acid and carbamazepine combination, or felbamate (felbatol).



WO 02/03915 A2

METHODS OF PRODUCING WEIGHT LOSS AND TREATMENT OF OBESITY

FIELD OF THE INVENTION

The present invention relates to pharmaceutical compositions, whose mechanisms of action(s) are at the kainate/AMPA receptor that can be used to treat the medical condition of obesity.

BACKGROUND OF THE INVENTION

Obesity is one of the most common medical disorders, which affects 30-40 % of the population of which 10% may be severe and morbid. Complications of obesity include insulin resistance, diabetes mellitus, hypertension, cardiovascular disease, pseudotumor cerebri, hyperlipidemia, sleep apnea, cancer, pulmonary hypertension, cholecystitis, and osteoarthritis. The mortality from obesity is estimated at 300,000 to 400,000 per annum in the United States. Obesity in humans is commonly measured by the BMI (body mass index) which is the weight in kilograms divided by the height in meters squared. The degree of obesity is determined by comparisons against standard deviations above the means for males and females. The exact etiology of obesity is unknown but occurs when energy intake exceeds energy expenditure. The amount and distribution of body fat may have some genetic predisposition and be under some hormonal control. Hypothalamic structures, which have complex interconnections with the limbic system and other brain structures, control appetite. Some neurochemicals known to be involved in appetite control include: leptin, a substance released from adipose tissue, GLP-1 (glucagon-like peptide1) which promotes satiety, and neuropeptide-Y, a potent stimulator of appetite.

SUMMARY OF THE INVENTION

The present invention proposes a theory of obesity in which dysfunction of the AMPA/kainate and/or NMDA is a contributing etiology. Administration of drugs whose mechanism or action is antagonism of the AMPA/kainate receptor, with or without the combination of glycine-site antagonists, is proposed as a treatment for obesity.

DETAILED DESCRIPTION OF CERTAIN PREFERRED EMBODIMENTS

Excitatory amino acids are divided into NMDA and non-NMDA [kainate (KA) and AMPA] subtypes. NMDA, KA, and AMPA receptors are concentrated in the lateral hypothalamus, hippocampus and cerebral cortex.

The NMDA receptor complex is located on the neuronal cell surface and is comprised of multiple binding sites as well as an ion-channel with several internal binding sites. Unique features of the NMDA receptor include strychnine insensitivity, blockade by physiological concentrations of Mg^{++} , a high permeability to Ca^{++} , and coactivation by glycine. However, the interaction of the NMDA, KA, and AMPA receptors are not completely understood. The release of endogenous glycine, which acts at the glycine site of the NMDA receptor, from the striatal neurons in primary culture, has been reported to be regulated principally by excitatory amino acids at the kainate receptor system.

High densities of NMDA receptors located in the ventromedial (VMH), medial (MH) and lateral hypothalamus (LH) suggest a prominent role in neuroendocrine and autonomic regulation. The NMDA receptor appears to play a significant physiological role in natural eating and body weight regulation. Studies suggest that NMDA, AMPA, and KA receptor subtypes exist on LH neurons and their activation

triggers intense eating. In animals, NMDA receptor blockade of the LH caused marked body weight loss and reduced eating at the onset of the circadian feeding period and after fasting.

5 Glutamate and its agonists act within the LH to elicit eating and thus participate of the LH in the regulation of eating behavior. LH injections of either glutamate, KA, AMPA, and NMDA elicit a strong dose-dependent eating response. Lateral hypothalamic NMDA blockade reduced the
10 eating response to NMDA and suppressed both deprivation-induced eating and nocturnal eating. These results suggest that endogenous LH glutamate and the NMDA receptor are both involved in the regulation of natural eating, circadian factors, and eating behavior. Thus, NMDA receptors in the
15 LH may participate in the long-term regulation of food intake and body weight control. There is evidence that they also provide mechanisms to account for the experience-dependent plasticity and modulation by motivational state exhibited by some feeding-related LH neurons.

20 Feeding and drinking are impaired in rats previously given intrahypothalamic injections of kainic acid, a neurotoxic agent that destroys neurons with cell bodies in the lateral hypothalamus but spares fibers of passage. Kainic acid (KA), D,L- α -amino-3-hydroxy-5-methyl-isoxazole
25 proprionic acid (AMPA) or N-methyl-D-aspartate (NMDA) injected into the lateral hypothalamus each elicited eating in a dose-dependent fashion. The effectiveness of all three of these agonists suggests that these receptor subtypes are each expressed by some LH neurons that
30 regulate eating. There is substantial hypothalamic NMDA and AMPA receptor subtype gene expression and evidence that

both NMDA and non-NMDA receptors are critical for eating behavior and weight control. NMDA receptors may mediate neural plasticity and learning, which may contribute to the pathophysiology of obesity. Thus, glutamate and several of its agonists can act within the LH to elicit eating, suggest that glutamate and several of its receptor subtypes may participate in LH regulation of eating behavior.

Glutamate and its agonists are relatively ineffective in the PVN (paraventricular nucleus) and PFH (perifornical hypothalamus). These hypothalamic sites are involved in the regulation of eating behavior and in mediating some of glutamate's other physiological effects. Specifically, the PVN has been identified as a primary locus for the feeding-stimulatory effects of norepinephrine, galanin and γ -aminobutyric acid, while the PFH has been identified as a primary site for the feeding-stimulatory effects of neuropeptide Y and the feeding-suppressive effects of dopamine and epinephrine. Moreover, both sites appear to express functional glutamate-sensitive receptors, since PVN or PHF injections of glutamate can produce autonomic and endocrine effects. That glutamate acts primarily in the LH, and neuropeptide γ in the PFH to elicit eating, suggests that these adjacent sites are functionally distinct with respect to eating. Some LH eating-regulatory neurons express multiple excitatory amino acid receptors, possibly including the kainate, AMPA, and NMDA subtypes. Autoradiographic studies reveal low levels of hypothalamic NMDA and AMPA binding, and only moderate levels of kainate binding. In contrast, there is evidence for substantial hypothalamic NMDA and AMPA receptor mediation of other physiologic effects in the hypothalamus.

In the lateral hypothalamus, glutamate may act as a physiological mediator of feeding behavior since injection of glutamate elicited eating in satiated rats. The application of AMPA, KA, and NMDA to the lateral hypothalamus was able to trigger intense eating. The NMDA receptor gates Ca^{++} , is both ligand and voltage sensitive, and is regulated by multiple modulatory sites. These mechanisms may account for some of the experience-dependent plasticity and modulation by motivational state exhibited by some feeding-related LH neurons.

To elucidate the role of the glycine binding site on the NMDA receptor in feeding control, a glycine site-antagonist injected into the LH of satiated rats blocked the stimulatory eating response of NMDA. Glycine-site antagonism also blocked the stimulatory feeding effects of KA and AMPA in a dose-dependent manner. In addition, bilateral LH injection of a glycine site antagonist also suppressed eating produced by fasting. These findings suggest an important role of the NMDA receptor glycine site in LH regulation of eating behavior.

Thus the glycine site antagonist 7-chlorokynurenic acid (7-CK) was injected into the lateral hypothalamus (LH) of satiated rats before LH injection of NMDA. 7-CK blocked the 6- to 10-gram eating response elicited by NMDA. LH pretreatment with glycine, suggesting a specific action at the glycine site reversed this block. In contrast to the suppression produced by high doses, 7-CK at 0.1 nmol enhanced NMDA-elicited eating. In addition, 7-CK at a dose of 0.1 nmol suppressed feeding elicited by KA or AMPA, but at 10 nmol it suppressed eating elicited by AMPA while enhancing eating elicited by KA. Finally, bilateral LH injection of 7-CK effectively suppressed eating produced by

fasting.

Specifically, the 0.1-nmol dose of 7-CK that enhanced NMDA-elicited eating suppressed KA-elicited eating, and the 10-nmol dose that blocked NMDA-elicited eating markedly enhanced the KA-elicited response. 7-CK had a significant effect on eating elicited by AMPA, with both doses of 7-CK suppressed eating, with the 0.1-nmol dose virtually blocking it.

In a further study, D-AP5 reduced by 72-90% the 10-gram eating response elicited by NMDA without affecting the quantitatively similar eating responses elicited by kainic acid or AMPA. This treatment also suppressed deprivation-induced eating responses by as much as 61% and nocturnal eating by as much as 40%. The injection of D-AP5 into the LH bilaterally for 8n days caused a 65% reduction in daily food intake and body weight loss of up to 13 grams/day. These findings suggested that endogenous LH glutamate acts to regulate natural eating and body weight and that NMDA receptors participate in these functions. Endogenous glutamate and NMDA receptors are important modulators of natural eating, specifically that generated by nutrient deprivation and circadian factors. Many LH neurons are also sensitive to food-related taste, olfactory, auditory, and visual inputs and NMDA receptors may participate in mediating the effects on eating behavior of some of these inputs.

Conversely, other studies have reported that the strychnine-insensitive glycine-binding site antagonist, 7-chlorokynurenic acid (7CK) dose-dependently increased food intake in rats when administered into the intracerebral ventricles or paraventricular nucleus of the hypothalamus. In addition, D-serine, a strychnine-insensitive glycine-

binding site agonist, dose-dependently antagonized 7CK induced feeding. Other brain structures may also be involved in appetite control since median raphe injection of excitatory amino acid antagonists elicited eating.

5 I believe that one etiology of obesity is dysfunction of the NMDA/AMPA/kainate receptors. This may include abnormal quantities of receptors, abnormal function of receptors, or abnormal interaction of these receptors.

Topomax

10 Topiramate (topomax) is a sulfamate-substituted monosaccharide, which is approved as adjunctive therapy in patients with partial onset seizures, or primary generalized tonic-clonic seizures. Topiramate (TPM) has the molecular formula $C_{12}H_{21}NO_8S$ and a molecular weight of 339.37 and is designated chemically as 2,3:4,5-DI-O-isolpropylidene-B-D-fructopyranose sulfamate. Tablets are 15 available as 25 mg, 100 mg, and 200 mg round tablets for oral administration.

The known mechanisms of action of TPM include: a 20 voltage-sensitive sodium channel blocking action, potentiation of the effects of GABA, and antagonizing the kainate/AMPA (α -amino-3-hydroxy-5-methylisoxazole-4-proprionic acid; non-NMDA) subtype of excitatory acid (glutamate) receptor. Using magnetic resonance 25 spectroscopy, TPM was found to acutely increase brain levels of cerebral GABA within three hours of oral administration. TPM has no effect on the activity of N-methyl-D-aspartate (NMDA) at the NMDA receptor subtype. The effects of TPM are concentration-dependent within the 30 range of 1 μM to 200 μM . TPM also inhibits some isoenzymes of carbonic anhydrase (CA-II and CA-IV) but this

pharmacological effect is weaker than that of acetazolamide, a known carbonic anhydrase inhibitor, and is currently not thought to be a major factor in the anti-epileptic activity of TPM.

5 Absorption of TPM is rapid, with time to peak (TTP) plasma concentrations occurring about 2 hours following a 400 mg dose. The relative bioavailability of TPM from the tablet formation is not affected by food. The pharmacokinetics of TPM is linear with dose proportional
10 increases in plasma concentration over the dose range of 200 to 800 mg/day. The mean plasma concentration half-life is 21 hours after either single or multiple doses. Steady state is reached in about 4 days in patients with normal renal function. TPM is 13-17% bound to human plasma
15 proteins over the concentration range of 1-250 µg/ml. The mean TTP plasma concentration was between 1.4 and 4.3 hours with TPM doses ranging from 100 to 1200 mg. Absorption is nearly complete, with 81-95% of a 100-mg dose recovered in urine. TPM is a lipophilic substance with a volume of
20 distribution of 0.80-0.55 L/kg and plasma protein binding is minimal, approximately 15%. Elimination of TPM (and its metabolites) is predominantly renal, with 50-80% of a dose excreted as unchanged TPM. TPM is not extensively
25 metabolized and is primarily eliminated unchanged in the urine (approximately 70% of an administered dose). Six metabolites have been identified in humans, none of which constitutes more than 5% of an administered dose. The metabolites are formed via hydroxylation, hydrolysis, and glucuronidation. There is evidence of renal tubular
30 reabsorption of TPM. In humans, plasma clearance (CL/F) is approximately 20 to 30 mL/min following oral

administration.

In patients with normal renal function, the elimination half-life for TPM is 20-30 hours, allowing steady-state plasma concentrations to be achieved in 4-8
5 days. In patients with moderate or severe renal impairment, TPM plasma levels are increased as plasma and renal clearances are reduced. In patients with hepatic impairment, TPM plasma concentrations increased by 29%. The clearance of TPM was reduced by 42% in moderate renal
10 impairment (creatinine clearance 30-69 mL/min/1.73m²) and by 54% in severe renal impairment (creatinine clearance <30 mL/min/1.73m²) compared to normal renal function subjects (creatinine clearance >30 mL/min/1.73m²). In hepatically impaired subjects, the clearance of TPM may be decreased.
15 However, the mechanism(s) underlying the decrease is not well understood.

The recommended titration of TPM therapy is an initiation at 50 mg/day with weekly increases of 50 mg/day. The current recommended total daily dose of TPM as
20 adjunctive epilepsy therapy is 400 mg in two divided doses. In a monotherapy study, adult patients were randomized to a treatment dosage of 1000 mg/day administered twice daily and were maintained at this dose for 112 days. In some studies, doses of 1600 mg/day and long-term administration
25 (>60 months) have been achieved, with mean doses of TPM averaging 700 mg/day. A therapeutic plasma concentration range has not been established for TPM and specific plasma concentrations were not correlated with the occurrence of adverse events. Therefore, clinicians have been advised to
30 use clinical effect as a guide for titrating TPM.

Common adverse events of TPM at dosages of 200 to 400 mg/day in controlled trials in adults with partial onset

seizures, primarily generalized tonic-clonic seizures, or Lennox-Gastaut syndrome were: somnolence, dizziness, ataxia, speech disorders and related speech problems, psychomotor slowing, abnormal vision, difficulty with memory, paresthesia and diplopia. These adverse events occurred at a greater frequency in TPM-treated patients and did not appear to be dose-related.

Conversely, the most common dose-related adverse events at dosages of 200 to 1,000 mg/day were: fatigue, nervousness, difficulty with concentration or attention, confusion, depression, anorexia, language problems, anxiety, mood problems, and weight decrease. The most common dose-related adverse events with dosages of 200-1000 mg/day were fatigue, nervousness, difficulty with concentration or attention, confusion, and depression. These adverse events have been attributed to the rapid rate of dose titration of 200 mg/day. However, the majority of these adverse events abated after 4 months of treatment. In a study of the acute and steady-state cognitive effects of TPM on young healthy adults, statistically significant declines on measures of attention and word fluency at both acute doses and at both 2- and 4- week test periods.

In clinical trials on seizure patients, weight loss averaged 2-8 kg (2-8% decline from baseline weight). Patients experiencing weight loss were more likely to report anorexia or loss of appetite. Paradoxically, weight loss tended to be higher in patients also being treated with valproate, or valproate with carbamazepine. Both of these drugs have been associated with weight gain when used as monotherapy. Weight loss generally peaked after 12-15 months of therapy, with body weight gradually returning toward pre-topiramate levels after prolonged therapy. As

adjunctive therapy in adult seizure patients, weight decrease occurred more frequently: control 3%, 200-400 mg/day of TPM 9%, and 600-1,000 mg/day of TPM 13%. In a monotherapy study in epilepsy patients, anorexia occurred at 100 mg (13%) and 1000 mg (42%) doses. In another study in epileptic patients, weight loss occurred in 5/12 patients (12-42 pounds) while weight increase was observed in 2/12 patients (12-25 pounds). In a long-term study of epileptic patients receiving TPM, 26% of patients treated for up to 2.2 years experienced weight loss.

Central nervous system adverse events are associated with the use of TPM. These included psychomotor slowing, difficulty with concentration, and speech or language problems, in particular, word-finding difficulty and somnolence and/or fatigue. Additional nonspecific CNS effects occasionally observed with TPM as add-on therapy include dizziness or imbalance, confusion, memory problems, and exacerbation of mood disturbances (i.e., irritability and depression). Reports of psychomotor slowing, speech and language problems, and difficulty with concentration and attention were common in adults. Somnolence and fatigue were the most frequently adverse event during the clinical trials, which were generally mild to moderate and occurred early in therapy. While the incidence of somnolence does not appear to be dose-related, that of fatigue increases at dosages above 400 mg/day.

Adverse events include renal calculi (kidney stones) which occurred in 1.5% of adults, an incidence about 2-4 times that expected in a similar, untreated population. This may be due to the fact that TPM is a weak carbonic anhydrase inhibitor and promote stone formation by reducing urinary citrate excretion and by increasing urinary pH.

The concomitant use of TPM in patients on a ketogenic diet may create a physiological environment that increases the risk of kidney stone formation, and is therefore not recommended.

5 In controlled clinical trials, 11% of patients receiving 200 to 400 mg/day as adjunctive therapy discontinued due to adverse events and increased at dosages above 400 mg/day. Adverse events associated with discontinuing therapy included somnolence, dizziness,
10 anxiety, difficulty with concentration or attention, fatigue, and paresthesia and increased at dosages above 400 mg/day. In post-marketing data, reported adverse experiences included hepatic failure, hepatitis, pancreatitis, and renal tubular acidosis.

15 The efficacy of TPM in non-epileptic patients with obesity is unknown. There is currently no published study on the effects of TPM in patients with obesity. In addition, the possibility exists that cognitive impairment in patients receiving TPM may prevent adequate dose
20 administration to produce weight loss in obese patients.

Lamotrigene

Lamotrigene (LTG) is an anticonvulsant whose mechanism of action is blocking the sodium channel. A common adverse event is anorexia and nausea (6-15%) which occurred at both
25 doses of 100 mg and 200 mg, adjunctive therapy, and monotherapy. LTG had no significant effect on body weight when administered for 12 months or up to 144 weeks. The combination of LTG and TPM in obese patients may be additive in their effects by producing both anorexia and
30 possibly weight loss.

Felbamate

Felbamate (FBM) was approved as an anti-epileptic drug

in 1993. FBM functions as a glycine site NMDA antagonist but other mechanisms of action reported include interaction at the kainate/AMPA receptor, modulation of the Na⁺⁺ channel conductance, facilitation of the GABA receptor, an
5 interaction at the L-type calcium channels.

In clinical studies prior to FDA approval, FBM was evaluated as add-on therapy (in combination with other standard epileptic drugs) in patients with Lennox-Gestalt Syndrome (intractable seizures with mental retardation).
10 Several mild cases of thrombocytopenia and neutropenia were reported but these were reversible and common adverse events of the other concomitant antiepileptic drugs. Prior to drug approval, there were no reported cases of aplastic anemia or hepatic dysfunction, when standard drug
15 monitoring parameters were performed. Drug interactions included elevation of dilantin, depakote, phenobarbitol levels and the toxic epoxide metabolite of tegretol. In 1994, there were reports of FBM causing aplastic anemia and hepatic failure. As a result, the FDA and manufacturer
20 sent a letter to all physicians in August 1994 requiring the withdrawal of FBM from patients whom were considered at risk for aplastic anemia. Warnings were added to the prescribing information and both hematological and hepatic indices were required every two weeks. In a study on
25 epileptic patients published after the FDA warning, 111 patients with refractory epilepsy were treated with adjunctive FBM therapy. No cases of aplastic anemia or hepatic dysfunction were observed when serum levels of all anti-convulsant drugs were monitored. Recent analysis has
30 suggested that only several cases of aplastic anemia could be possibly be attributed to FBM monotherapy. In the absence of any adverse event, recent analysis has suggested

that FBM, even in adjunctive therapy, is relatively safe after usage for at least one year.

FBM has been known to induce anorexia and weight loss in patients treated for epilepsy. In a study of FBM as
5 add-on therapy in partial epilepsy in non-obese patients, weight loss was transient and returned to baseline after twenty weeks. Weight loss of 5% was reported in non-obese patients of FBM monotherapy. The weight loss in these
10 patients was attributed to nausea and vomiting as well as the withdrawal of other medications whose side effects were weight gain. We postulate that weight loss from FBM is most likely due to NMDA receptor and non-NMDA modulation of the various hypothalamic structures and receptors involved in appetite control. The mechanisms of action and
15 anecdotal reports of weight loss with monotherapy suggest that FBM is a potentially excellent therapeutic agent in subjects with obesity. In a study of cognitive effects of FBM and tegretol in epileptic patients, FBM was found to increase both cognition and fine motor function. In
20 patients who suffer cognitive decline while receiving TPM, adjunctive FBM may be able to reverse these abnormal cognitive adverse effects.

I claim:

1. A method for treating a human suffering from obesity, comprising the steps of:

administering to the human a compound whose
5 mechanism of action includes antagonism of the kainite and/or AMPA receptor.

2. The method of claim 1, further comprising administering to said human topiramate monotherapy at dose ranges of 1 mg to 5000 mg per day.

10 3. The method of claim 1, further comprising administering to said human monotherapy at dose ranges more preferably from 400-2000 per day.

4. The method of claim 1, further comprising administering to said human topiramate in combination with
15 lamotrigine.

5. The method of claim 1, further comprising administering to said human topiramate in combination with valproic acid.

6. The method of claim 1, further comprising
20 administering to said human topiramate in combination with both valproic acid and carbamazepine.

7. The method of claim 1, further comprising administering to said human topiramate in combination with felbamate.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 January 2002 (17.01.2002)

PCT

(10) International Publication Number
WO 02/03915 A3

(51) International Patent Classification⁷: **A61K 31/70**,
31/55, 31/22, 31/19

(21) International Application Number: PCT/US01/21509

(22) International Filing Date: 9 July 2001 (09.07.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/613,692 10 July 2000 (10.07.2000) US

(71) Applicant and

(72) Inventor: **KOZACHUK, Walter, E.** [US/US]; 11403-B
Cam Court, Kensington, MD 20895-1313 (US).

(74) Agent: **LINIAK, Thomas, P.**; Liniak, Berenato, Longacre & White, LLC, Suite 240, 6550 Rock Spring Drive, Bethesda, MD 20817 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
23 May 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHODS OF PRODUCING WEIGHT LOSS AND TREATMENT OF OBESITY

(57) Abstract: Methods are disclosed for the acute and chronic treatment of obesity using drugs whose mechanism includes the interaction and antagonism of the kainate/AMPA receptor. Methods are disclosed for employing the drug topiramate (topomax) as monotherapy or in combination therapy with lamotrigine, valproic acid, valproic acid and carbamazepine combination, or felbamate (felbatol).

WO 02/03915 A3

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US01/21509

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/70, 31/55, 31/22, 31/19

US CL : 514/23, 217, 551, 557

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/23, 217, 551, 557

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,753,693 A (SHANK) 19 May 1998 (19.05.98), see the entire document.	1-7
A	US 5,753,694 A (SHANK) 19 May 1998 (19.05.98), see the entire document.	1-7



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"G" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

08 SEPTEMBER 2001

Date of mailing of the international search report

16 JAN 2002

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

RAYMOND J. HENLEY III

Telephone No. (703) 308-1235